



# Introduction to Mathematics and Modeling

## lecture 5

### The chain rule and optimization

**UNIVERSITY OF TWENTE.**

academic year : 18-19

lecture : 5

build : December 10, 2018


slides : 31

This week

[intro](#)



- 1 Section 3.6: the chain rule
- 2 Section 3.8: derivatives of logarithms (only pages 176–181)
- 3 Section 4.1: extreme values




- The **composition of  $f$  and  $g$**  is the function that maps  $x$  to  $f(g(x))$
- The composition is denoted as  $f \circ g$ , and is pronounced as “ $f$  after  $g$ ”.
- Example: let  $f(x) = x^2$  and let  $g(x) = x + 1$ , then

$$f \circ g(x) = f(g(x)) = f(x + 1) = (x + 1)^2 = x^2 + 2x + 1$$

and

$$g \circ f(x) = g(f(x)) = g(x^2) = x^2 + 1.$$

 In general  $f \circ g$  and  $g \circ f$  are *not* identical.

## Composition with a linear function

- Let  $f(x) = ax + b$  and  $g(x) = \sin(x)$  and define  $h = f \circ g$ , then

$$h(x) = f \circ g(x) = f(g(x)) = a \sin(x) + b$$


- Using the sum rule and constant multiple rule we know that

$$h'(x) = a \cos(x)$$

- Now let  $h = g \circ f$  then

$$h(x) = g(f(x)) = \sin(ax + b)$$

The sum- and constant multiple rule cannot be applied



- Consider the special case  $h(x) = \sin(3x)$ . The graph of  $h$  is obtained by scaling  $\sin x$  in horizontal direction.
- The slopes of all tangents are scaled too!
- By scaling back  $\sin(3x)$  in vertical direction, this effect is cancelled out:

$$\frac{d}{dx} \left( \frac{1}{3} \sin(3x) \right) = \cos(3x),$$

in other words:  $\frac{d}{dx} \sin(3x) = 3 \cos(3x)$ .

- We see that

$$f(x) = \sin(ax) \quad \Rightarrow \quad f'(x) = a \cos(ax)$$

- By shifting a graph horizontally, the slopes must shift accordingly:

$$f(x) = \sin(ax + b) \quad \Rightarrow \quad f'(x) = a \cos(ax + b)$$

### Chain rule, simple version

Let  $f$  be a differentiable function. Then for any constant  $a$  and  $b$  the following holds:

$$\frac{d}{dx} (f(ax + b)) = af'(ax + b).$$

**⚠ Warning:**  $\frac{d}{dx} (f(ax + b))$  is the derivative of the composition  $f(ax + b)$ , but  $f'(ax + b)$  is the composition of  $f'$  and  $y = ax + b$ .

- The derivative of  $\sin(2x)$  is  $2 \cos(2x)$ .

- Define  $y = \sqrt{5 - 3x}$ , then

$$\frac{dy}{dx} = -\frac{3}{2\sqrt{5-3x}}$$

since  $\frac{d}{dx}(\sqrt{x}) = \frac{1}{2\sqrt{x}}$ .

Also: write  $5 - 3x = (-3)x + 5$ , hence  $a = -3$  and  $b = 5$ .

- $\frac{d}{dx} \left( \frac{1}{2e^x} \right) =$

## Application: the derivative of exponential functions

- See lecture 4: if we define  $f(x) = a^x$ , then

$$f'(x) = k_a a^x$$

where

$$k_a = \lim_{h \rightarrow 0} \frac{a^h - 1}{h} = f'(0).$$

- With the simple version of the chain rule we can prove:

$$k_a = \ln a$$

$$\frac{d}{dx} (a^x) =$$

**Chain rule**

Let  $f$  and  $g$  be differentiable functions, then



$$\frac{d}{dx}f(g(x)) = f'(g(x))g'(x).$$

- In words: multiply the composition of the derivative of  $f$  with  $g$  by the derivative of  $g$ .
- Work inward:
  - differentiate the 'outer function'  $f$ , but keep the 'inner function'  $g$  intact;
  - then multiply with the derivative of the 'inner function'  $g$ .


**Example**

Find the derivative of  $h(x) = (3x^2 + 1)^2$ .

- The function  $h$  is equal to  $h = f \circ g$ , where
$$f(x) = x^2 \quad \text{and} \quad g(x) = 3x^2 + 1.$$
- Apply the chain rule:
$$h'(x) =$$



- If a function is named  $f$ , the derivative is denoted as  $f'$ .
- If  $y$  is an anonymous function of  $x$ , the derivative is denoted as  $\frac{dy}{dx}$ .



If  $y$  is a function of  $x$  and  $z$  is a function of  $y$ , then  $z$  is (by composition) a function of  $x$ . In this case the chain rule is

$$\frac{dz}{dx} = \frac{dz}{dy} \frac{dy}{dx}.$$

- ⚠ Note that  $\frac{dz}{dy}$  is expressed in terms of  $y$ , hence afterwards you should replace all occurrences of  $y$  with the corresponding expression in  $x$ .

### Example

Let  $y = 3x^2 + 1$  and  $z = y^2$ , find  $\frac{dz}{dx}$ .

- Apply the chain rule (anonymous variant):

$$\frac{dz}{dx} =$$

**Example**

Find the derivative of  $f(x) = \frac{1}{\sqrt{x^2 + 1}}$ .

- Avoid using the quotient rule by writing

$$f(x) = (x^2 + 1)^{-1/2}.$$

- Apply the chain rule:

$$f'(x) =$$

**Example**

Calculate the derivative of  $f(x) = \sqrt{\frac{1-x^2}{1+x^2}}$ .

- Combine the chain rule with the quotient rule:

$$f'(x) =$$

- The **logarithmic function with base  $a$**  is the inverse of the base- $a$  exponential function:

$$y = a^x \iff x = \log_a y$$

- The **natural logarithm** is the logarithm with base  $e$ :

$$\ln x = \log_e x$$

where

$$e \approx 2.71828\dots$$

- Examples:

$$\log_2 8 = 3 \quad \text{because} \quad 2^3 = 8$$

$$\log_{10} 100 = 2 \quad \text{because} \quad 10^2 = 100$$

$$\ln e\sqrt{e} = \frac{3}{2} \quad \text{because} \quad e^{\frac{3}{2}} = e\sqrt{e}$$

$$\log_a 1 = 0 \quad \text{and} \quad \log_a a = 1$$

$$\log_a xy = \log_a x + \log_a y$$

$$\log_a \frac{x}{y} = \log_a x - \log_a y$$

$$\log_a \frac{1}{y} = -\log_a y$$

$$\log_a x^p = p \log_a x$$

$$\log_a x = \frac{\log_b x}{\log_b a}, \quad \text{in particular} \quad \log_a x = \frac{\ln x}{\ln a}$$

$$a^x = b^{x \log_b a}, \quad \text{in particular} \quad a^x = e^{x \ln a}$$



- Note that  $e^x$  and  $\ln(x)$  are each others inverse:

$$e^{\ln(x)} = x.$$


- Now take derivatives on both sides and apply the chain rule to the left-hand side:

$$e^{\ln(x)} \ln'(x) = 1,$$

$$x \ln'(x) = 1,$$

$$\ln'(x) = \frac{1}{x}.$$

⚠ This holds for  $x > 0$ .



### Theorem

The derivative of  $\log_a x$  is  $\frac{1}{x \ln(a)}$ .

- From the change-of-base formula for logarithms follows

$$\log_a(x) = \frac{\ln x}{\ln a}.$$

- Apply the constant-multiple rule:

$$f'(x) =$$


**Example**

Find the derivative of  $f(x) = \ln(x^2 + 3)$ .

- Apply the chain rule:

$$f'(x) =$$

## Extreme values of a function



Consider a function  $f: D \rightarrow \mathbb{R}$ .

- $f$  has an **absolute maximum** in  $c$  if


$$f(x) \leq f(c) \quad \text{for all } x \in D$$

- $f$  has an **absolute minimum** in  $c$  if

$$f(x) \geq f(c) \quad \text{for all } x \in D$$

⚠ Extreme values do not necessarily have to exist!


?? If they exist, how do we find them?



On  $D = (-\infty, \infty)$  the function  $f(x) = x^2$  has

- an absolute minimum in  $x = 0$ ;
- no absolute maximum.


The range of  $f$  is  $[0, \infty)$ .



On  $D = [0, 2]$  the function  $f(x) = x^2$  has

- an absolute minimum in  $x = 0$ ;
- an absolute maximum in  $x = 2$ .


The range of  $f$  is  $[0, 4]$ .



On  $D = (0, 2]$  the function  $f(x) = x^2$  has

- no absolute minimum;
- an absolute maximum in  $x = 2$ .


The range of  $f$  is  $(0, 4]$ .



On  $D = (0, 2)$  the function  $f(x) = x^2$  has

- no absolute minimum;
- no absolute maximum.

The range of  $f$  is  $(0, 4)$ .




**Extreme Value Theorem**






Theorem 1, page 223

A **continuous function** on a **finite closed interval** attains both an **absolute maximum** and an **absolute minimum**.

 The theorem tells us that extreme values do exist, but *not* where to find them!


Local extremes



**Definition**

Consider a function  $f: D \rightarrow \mathbb{R}$ .

- $f$  has a **local maximum** in  $c$  if
 
$$f(x) \leq f(c) \quad \text{for all } x \text{ in an environment of } c$$
- $f$  has a **local minimum** in  $c$  if
 
$$f(x) \geq f(c) \quad \text{for all } x \text{ in an environment of } c$$



**First Derivative Theorem**

Theorem 2, page 225


If  $f$  is differentiable at  $c$  and  $f$  attains a local maximum or local minimum at  $c$  then  $f'(c) = 0$ .

**Definition**

The number  $c$  is a **critical point of  $f$**  if

$$f'(c) = 0 \quad \text{or} \quad f \text{ is not differentiable at } c.$$

Critical points are *candidates* for a local maximum or minimum.



The function  $F$  is defined on the domain  $[a, b]$ .

- Critical points:  $c_1, c_2, c_3, c_4$  and  $c_5$ .
- Local extremes:  $a, c_2, c_3, c_4$  and  $b$ .
- Absolute extremes:  $a$  and  $c_4$ .

Recipe for computing the extreme values of a continuous function

$$f: [a, b] \rightarrow \mathbb{R}$$

- 1 Find *all* critical points of  $f$  in  $[a, b]$ , i.e., solve the equation  $f'(x) = 0$  and retain all solutions  $x$  in  $[a, b]$ ; then add all points where  $f$  is not differentiable.
- 2 Evaluate  $f$  at the critical points and at the end points  $x = a$  and  $x = b$ .
- 3 Take the largest and smallest values found in step 2: these are the absolute maximum and minimum of  $f$  on the interval  $[a, b]$ .

**Example**

Find extremes for  $f(x) = 10x(2 - \ln x)$  on  $[1, e^2]$ .

- 1 Find the critical points:
- 2 Evaluate  $f$  at the critical points and at the endpoints:

$$f(1) = \qquad f(e^2) =$$

- 3 Take the largest and smallest values of step 2:
  - The absolute maximum is
  - The absolute minimum is

**Example**

Find extremes for  $f(x) = xe^{-x}$  on  $[-1, 1]$ .

**1** Find the critical points:

**2** Evaluate  $f$  at the critical points and at the endpoints:

$$f(-1) =$$

$$f(1) =$$

**3** Take the largest and smallest values of step 2:

- The absolute maximum is
- The absolute minimum is

**Example**

Find extremes for  $f(x) = 3x^2 - 2x^3$  on  $[-\frac{1}{2}, 2]$ .

**1** Find the critical points:

**2** Evaluate  $f$  at the critical points and at the endpoints:

$$f\left(-\frac{1}{2}\right) =$$

$$f(2) =$$

**3** Take the largest and smallest values of step 2:

- The absolute maximum is
- The absolute minimum is